

Ausgewählte Publikationen der letzten 5 Jahre

Airik, R., Trowe, M.-O., Foik, A., Farin, H.F., Petry, M., Schuster-Gossler, K., Schweizer, M., **Scherer, G.**, Kist, R., and Kispert, A.: Hydroureteronephrosis due to loss of Sox9-regulated smooth muscle cell differentiation of the ureteric mesenchyme. *Hum. Mol. Genet.* **19**, 4918-4929 (2010).

Lardenois, A., Chalmel, F., Barrionuevo, F., Demougin, P., **Scherer, G.**, and Primig, M.: Profiling spermatogenic failure in adult testes bearing Sox9-deficient Sertoli cells identifies genes involved in feminization, inflammation and stress. *Reprod. Biol. Endocrinol.* **8**, 154 (2010).

Kim, G.-J., Georg, I., Scherthan, H., Merckenschlager, M., Guillou, F., **Scherer, G.**, and Barrionuevo, F.: Dicer is required for Sertoli cell function and survival. *Int. J. Dev. Biol.* **54**, 867-875 (2010).

Georg, I., Bagheri-Fam, S., Knowler, K.C., Wieacker, P., **Scherer, G.**, and Harley, V.: Mutations of the SRY-responsive enhancer of SOX9 are uncommon in XY gonadal dysgenesis. *Sex. Dev.* **4**, 321-325 (2010).

Barrionuevo, F., Georg, I., Scherthan, H., Lecureuil, C., Guillou, F., Wegner, M., and **Scherer, G.**: Testis cord differentiation after the sex determination stage is independent of Sox9 but fails in the combined absence of Sox9 and Sox8. *Dev. Biol.* **327**, 301-312 (2009).

Rastegar, S., Hess, I., Dickmeis, T., Nicod, J.C., Ertzer, R., Hadzhiev, Y., Thies, W.-G., **Scherer, G.**, and Strähle, U.: The words of the regulatory code are arranged in a variable manner in highly conserved notochord enhancers. *Dev. Biol.* **318**, 366-377 (2008).

Bagheri-Fam, S., Sim, H., Bernard, P., Jayakody, I., Taketo, M.M., **Scherer, G.**, and Harley, V.R.: Loss of Fgfr2 leads to partial XY sex reversal. *Dev. Biol.* **314**, 71-83 (2008).

Barrionuevo, F., Naumann, A., Bagheri-Fam, S., Taketo, M.M., **Scherer, G.**, and Neubüser, A.: Sox9 is required for invagination and survival of the otic placode in mice. *Dev. Biol.* **317**, 213-224 (2008).

Bastide, P., Darido, C., Pannequin, J., Kist, R., Robine, S., Marty-Double, C., Bibeau, F., **Scherer, G.**, Joubert, D., Hollande, F., Blache, P., and Jay, P.: Sox9 regulates cell proliferation and is required for Paneth cell differentiation in the intestinal epithelium. *J. Cell Biol.* **178**, 635-648 (2007).

Leipoldt, M., Erdel, M., Bien-Willner, G.A., Smyk, M., Theurl, M., Yatsenko, S., Lupski, J.R., Lane, A.H., Shanske, A.L., Stankiewicz, P., and **Scherer, G.**: Two novel translocation breakpoints upstream of SOX9 define borders of the proximal and distal breakpoint cluster region in campomelic dysplasia. *Clin. Genet.* **71**, 67-75 (2007).

Seymour, P.A., Freude, K.K., Tran, M.N., Mayes, E.E., Jensen, J., Kist, R., **Scherer, G.**, and Sander, M.: SOX9 is required for maintenance of the pancreatic progenitor cell pool. *Proc. Natl. Acad. Sci. USA* **104**, 1865-1870 (2007).

Temel, S.G., Gulten, T., Yakut, T., Saglam, H., Kilic, N., Bausch, E., Jin, W.J., Leipoldt, M., and **Scherer, G.**: Extended pedigree with multiple cases of XX sex reversal in the absence of SRY and of a mutation at the SOX9 locus. *Sex. Dev.* **1**, 24-34 (2007).

Barrionuevo, F., Bagheri-Fam, S., Klattig, J., Kist, R., Taketo, M.M., Englert, C., and **Scherer, G.**: Homozygous inactivation of Sox9 causes complete XY sex reversal in mice. *Biol. Reprod.* **74**, 195-201 (2006).