

## Main Publications

(Recent publications at PubMed)

- Metzger E, Imhof A, Patel D, Kahl P, Hoffmeyer K, Friedrichs N, Müller JM, Greschik H, Kirfel J, Ji S, Kunowska N, Beisenherz-Huss C, Günther T, Buettner R, Schüle R. (2010). Phosphorylation of histone H3T6 by PKCbeta(I) controls demethylation at histone H3K4. *Nature* 464, 792-6. PDF
- Metzger E, Yin N, Wissmann M, Kunowska N, Fischer K, Friedrichs N, Patnaik D, Higgins JM, Potier N, Scheidtmann KH, Buettner R, Schüle R. (2008). Phosphorylation of histone H3 at threonine 11 establishes a novel chromatin mark for transcriptional regulation. *Nat Cell Biol.* 10, 53-60. PDF
- Wissmann M, Yin N, Mueller JM, Greschik H, Fodor BD, Jenuwein T, Vogler C, Schneider R, Guenther T, Buettner R, Metzger E, Schuele R (2007). Co-operative demethylation by JMJD2C and LSD1 promotes androgen receptor-dependent gene expression. *Nat Cell Biol.* 9, 347-353. PDF
- Hublitz P, Kunowska N, Mayer UP, Muller JM, Heyne K, Yin N, Fritzsche C, Poli C, Miguet L, Schupp IW, van Grunsven LA, Potiers N, van Dorsselaer A, Metzger E, Roemer K, Schuele, R. (2005). NIR is a novel INHAT repressor that modulates the transcriptional activity of p53. *Genes Dev.* 19, 2912-24. PDF
- Metzger E, Wissmann M, Yin N, Muller JM, Schneider R, Peters AH, Gunther T, Buettner R, Schuele, R. (2005). LSD1 demethylates repressive histone marks to promote androgen-receptor-dependent transcription *Nature.* 437, 436-439. PDF
- Gunther T, Poli C, Muller JM, Catala-Lehnen P, Schinke T, Yin N, Vomstein S, Amling M, Schuele, R. (2005). Fhl2 deficiency results in osteopenia due to decreased activity of osteoblasts. *EMBO J.* 24, 3049-56. PDF
- Stehlin-Gaon C, Willmann D, Zeyer D, Sanglier S, Van Dorsselaer A, Renaud JP, Moras D, Schule R. (2003). All-trans retinoic acid is a ligand for the orphan nuclear receptor ROR beta. *Nat. Struct. Biol.* 10, 820-825. PDF
- Metzger E, Muller JM, Ferrari S, Buettner R, Schuele, R. (2003). A novel inducible transactivation domain in the androgen receptor: implications for PRK in prostate cancer. *EMBO J.* 15, 270-280. PDF
- Mueller, J.M., Metzger, E., Greschik, H., Bosserhoff, A-K., Mercep, L., Buettner, R. and Schuele, R. (2002). The transcriptional coactivator FHL2 transmits Rho signals from the cell membrane into the nucleus. *EMBO J.* 21, 736-748. PDF
- Stehlin C, Wurtz JM, Steinmetz A, Greiner E, Schuele R., Moras D, Renaud JP. (2001). X-ray structure of the orphan nuclear receptor RORbeta ligand-binding domain in the active conformation. *EMBO J.* 20, 5822-5831. PDF
- Mueller, J.M., Isele, U., Metzger, E., Rempel, A., Moser, M., Pscherer, A., Breyer, T., Holubarsch, C., Buettner, R. and Schuele, R. (2000) FHL2, a novel tissue-specific coactivator of the androgen receptor. *EMBO J.* 19, 359-369. PDF
- Schuele, R., Pundi, R., Kliewer, S., Ransone, L.J., Bolado, J., Yang, N., Verma, I.M. and Evans, R.M. (1990) Functional antagonism between oncoprotein c-Jun and the glucocorticoid receptor. *Cell* 62, 1217-1226.

Schuele, R., Umesono, K., Mangelsdorf, D., Bolado, J., Pike, W.J. and Evans, R.M. (1990). Jun/Fos and receptors for vitamin A and D recognize a common response element in the human osteocalcin gene. *Cell* 61, 497-504.

Schuele, R., Muller, M., Kaltschmidt, Ch. and Renkawitz, R. (1988) Many transcription factors interact synergistically with steroid receptors. *Science* 242, 1418-1420.

Schuele, R., Muller, M., Oztuka-Murakami, H. and Renkawitz, R. (1988) Cooperativity of the glucocorticoid receptor and the CACCC-box binding factor. *Nature* 322, 87-90.