

LITERATUR

- Behneke A, Behneke N, d'Hoedt B (2002): A 5-year longitudinal study of the clinical effectiveness of ITI solid-screw implants in the treatment of mandibular edentulism. *Int J Oral Maxillofac Implants* 17: 799-810
- Besimo C, Graber G (1994): A new concept of overdentures with telescope crowns on osseointegrated implants. *Int J Periodontics Restorative Dent* 14: 486-495
- Chugal N M, Clive J M, Spangberg L S (2001): A prognostic model for assessment of the outcome of endodontic treatment: Effect of biologic and diagnostic variables. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 91: 342-352
- Coca I, Lotzmann U, Poggeler R (2000): Long-term experience with telescopically retained overdentures (double crown technique). *Eur J Prosthodont Restor Dent* 8: 33-37
- Drago C J (2003): A retrospective comparison of two definitive impression techniques and their associated postinsertion adjustments in complete denture prosthodontics. *J Prosthodont* 12: 192-197
- Eisenburger M, Gray G, Tschernitschek H (2000): Long-term results of telescopic crown retained dentures--a retrospective study. *Eur J Prosthodont Restor Dent* 8: 87-91
- Ferrigno N, Laureti M, Fanali S, Grippaudo G (2002): A long-term follow-up study of non-submerged ITI implants in the treatment of totally edentulous jaws. Part I: Ten-year life table analysis of a prospective multicenter study with 1286 implants. *Clin Oral Implants Res* 13: 260-273
- Friedman S, Lost C, Zarrabian M, Trope M (1995): Evaluation of success and failure after endodontic therapy using a glass ionomer cement sealer. *J Endod* 21: 384-390
- Gastaldo J F, Cury P R, Sendyk W R (2004): Effect of the vertical and horizontal distances between adjacent implants and between a tooth and an implant on the incidence of interproximal papilla. *J Periodontol* 75: 1242-1246
- Gotfredsen K, Holm B (2000): Implant-supported mandibular overdentures retained with ball or bar attachments: a randomized prospective 5-year study. *Int J Prosthodont* 13: 125-130
- He J, Cen Y, Mao X (1997): [A three-dimensional finite element stress analysis of implant-supported prosthesis and its supporting tissue in the edentulous mandible. Part 6. The influence of superstructure in mandibular complete implant overdenture and its supporting tissue]. *Hua Xi Kou Qiang Yi Xue Za Zhi* 15: 67-69
- Heckmann S M, Schrott A, Graef F, Wichmann M G, Weber H P (2004): Mandibular two-implant telescopic overdentures. *Clin Oral Implants Res* 15: 560-569
- Heydecke G, Peters M C (2002): The restoration of endodontically treated, single-rooted teeth with cast or direct posts and cores: a systematic review. *J Prosthet Dent* 87: 380-386

- Hinode D, Tanabe S, Yokoyama M, Fujisawa K, Yamauchi E, Miyamoto Y (2006): Influence of smoking on osseointegrated implant failure: a meta-analysis. *Clin Oral Implants Res* 17: 473-478
- Jokstad A (2006): Implant retained or conventional dentures, which give more patients satisfaction? *Evid Based Dent* 7: 96-97
- Kerschbaum T (2004): Langzeitüberlebensdauer von Zahnersatz. Eine Übersicht. *Quintessenz* 55: 1113-1126
- Kiener P, Oetterli M, Mericske E, Mericske-Stern R (2001): Effectiveness of maxillary overdentures supported by implants: maintenance and prosthetic complications. *Int J Prosthodont* 14: 133-140
- Krennmair G, Weinlander M, Krainhofner M, Piehslinger E (2006): Implant-supported mandibular overdentures retained with ball or telescopic crown attachments: a 3-year prospective study. *Int J Prosthodont* 19: 164-170
- Kwakman J M, Voorsmit R A, Freihofer H P, Van Waas M A, Geertman M E (1998): Randomized prospective clinical trial of two implant systems for overdenture treatment: a comparison of the 2-year and 5-year results using the clinical implant performance scale. *Int J Oral Maxillofac Surg* 27: 94-98
- Löst C, Weiger R, Axmann-Krcmar D (1995): Prognose von Wurzelkanalbehandlungen unter Anwendung der lateralen Kondensationstechnik und eines Glasionomerzementsealers. *Dtsch Zahnärztl Z* 50: 897-901
- Mock F, Schrenker H, Stark H K (2005): Eine klinische Langzeitstudie von Teleskopprothesen. *Dtsch Zahnärztl Z* 60: 148-153
- Nitzan D, Mamlider A, Levin L, Schwartz-Arad D (2005): Impact of smoking on marginal bone loss. *Int J Oral Maxillofac Implants* 20: 605-609
- Pan S, Yin Y, Feng H (1999): Three-dimensional finite element analysis and comparison of stress distribution in overdentures supported with bar attachments and telescopic crowns. *Chin J Dent Res* 2: 21-30
- Petrie C S, Walker M P, Williams K (2005): A survey of U.S. prosthodontists and dental schools on the current materials and methods for final impressions for complete denture prosthodontics. *J Prosthodont* 14: 253-262
- Quiryneen M, Alsaadi G, Pauwels M, Haffajee A, van Steenberghe D, Naert I (2005): Microbiological and clinical outcomes and patient satisfaction for two treatment options in the edentulous lower jaw after 10 years of function. *Clin Oral Implants Res* 16: 277-287
- Rehmann P, Weber A, Wöstmann B, Fergner P (2007): Klinische Bewährung von Zähnen, die zur Verankerung einer Teilprothese mit Teleskopkronen versorgt wurden. *Dtsch Zahnärztl Z* 62: 99-103

- Schaffarzik S, Coca I (2001): Die Wurzelstiftkappe - häufig unterschätztes prothetisches Therapiemittel. *Zahn Parx* 4: 352-356
- Sjogren U, Hagglund B, Sundqvist G, Wing K (1990): Factors affecting the long-term results of endodontic treatment. *J Endod* 16: 498-504
- Smith D E, Toolson L B, Bolender C L, Lord J L (1979): One-step border molding of complete denture impressions using a polyether impression material. *J Prosthet Dent* 41: 347-351
- Stankiewicz N R, Wilson P R (2002): The ferrule effect: a literature review. *Int Endod J* 35: 575-581
- Stellingsma C, Vissink A, Meijer H J, Kuiper C, Raghoobar G M (2004): Implantology and the severely resorbed edentulous mandible. *Crit Rev Oral Biol Med* 15: 240-248
- Strub J R, Gaberthuel T W, Grunder U (1991): The role of attached gingiva in the health of peri-implant tissue in dogs. 1. Clinical findings. *Int J Periodontics Restorative Dent* 11: 317-333
- Tan H K, Hooper P M, Baergen C G (1996): Variability in the shape of maxillary vestibular impressions recorded with modeling plastic and a polyether impression material. *Int J Prosthodont* 9: 282-289
- Timmerman R, Stoker G T, Wismeijer D, Oosterveld P, Vermeeren J I, van Waas M A (2004): An eight-year follow-up to a randomized clinical trial of participant satisfaction with three types of mandibular implant-retained overdentures. *J Dent Res* 83: 630-633
- van Steenberghe D, Quirynen M, Naert I, Maffei G, Jacobs R (2001): Marginal bone loss around implants retaining hinging mandibular overdentures, at 4-, 8- and 12-years follow-up. *J Clin Periodontol* 28: 628-633
- Weischer T, Mohr C (2001): Implant-supported mandibular telescopic prostheses in oral cancer patients: an up to 9-year retrospective study. *Int J Prosthodont* 14: 329-334
- Wennstrom J, Lindhe J (1983): Role of attached gingiva for maintenance of periodontal health. Healing following excisional and grafting procedures in dogs. *J Clin Periodontol* 10: 206-221
- Wennstrom J L (1987): Lack of association between width of attached gingiva and development of soft tissue recession. A 5-year longitudinal study. *J Clin Periodontol* 14: 181-184
- Wenz H J, Hertrampf K, Lehmann K M (2001): Clinical longevity of removable partial dentures retained by telescopic crowns: outcome of the double crown with clearance fit. *Int J Prosthodont* 14: 207-213