

LITERATUR

Bergman B, Carlsson G E (1985): Clinical long-term study of complete denture wearers. *J Prosthet Dent* 53: 56-61

Dubojska A M, White G E, Pasiak S (1998): The importance of occlusal balance in the control of complete dentures. *Quintessence Int* 29: 389-394

Feine J S, Carlsson G E, Awad M A, Chehade A, Duncan W J, Gizani S, Head T, Lund J P, MacEntee M, Mericske-Stern R, Mojon P, Morais J, Naert I, Payne A G, Penrod J, Stoker G T, Tawse-Smith A, Taylor T D, Thomason J M, Thomson W M, Wismeijer D (2002): The McGill consensus statement on overdentures. Mandibular two-implant overdentures as first choice standard of care for edentulous patients. Montreal, Quebec, May 24-25, 2002. *Int J Oral Maxillofac Implants* 17: 601-602

Ferrigno N, Laureti M, Fanali S, Grippaudo G (2002): A long-term follow-up study of non-submerged ITI implants in the treatment of totally edentulous jaws. Part I: Ten-year life table analysis of a prospective multicenter study with 1286 implants. *Clin Oral Implants Res* 13: 260-273

Gausch K (1986): [Experiences with anterior/cuspid-controlled complete dentures]. *Dtsch Zahnarztl Z* 41: 1146-1149

Haraldson T, Karlsson U, Carlsson G E (1979): Bite force and oral function in complete denture wearers. *J Oral Rehabil* 6: 41-48

Heckmann S M, Schrott A, Graef F, Wichmann M G, Weber H P (2004): Mandibular two-implant telescopic overdentures. *Clin Oral Implants Res* 15: 560-569

Hofmann M, Knauer G (1990): [Studies on canine guidance in complete dentures]. *Dtsch Zahnarztl Z* 45: 566-570

Jokstad A (2006): Implant retained or conventional dentures, which give more patients satisfaction? *Evid Based Dent* 7: 96-97

Krennmair G, Weinlander M, Krainhofner M, Piehslinger E (2006): Implant-supported mandibular overdentures retained with ball or telescopic crown attachments: a 3-year prospective study. *Int J Prosthodont* 19: 164-170

McCord J F, Grant A A (2000): Identification of complete denture problems: a summary. *Br Dent J* 189: 128-134

Närhi T O, Hevinga M, Voorsmit R A, Kalk W (2001): Maxillary overdentures retained by splinted and unsplinted implants: a retrospective study. *Int J Oral Maxillofac Implants* 16: 259-266

Peroz I, Leuenberg A, Haustein I, Lange K P (2003): Comparison between balanced occlusion and canine guidance in complete denture wearers--a clinical, randomized trial. *Quintessence Int* 34: 607-612

- Rissin L, House J E, Manly R S, Kapur K K (1978): Clinical comparison of masticatory performance and electromyographic activity of patients with complete dentures, overdentures, and natural teeth. *J Prosthet Dent* 39: 508-511
- Rivera-Morales W C, Goldman B M (1997): Are speech-based techniques for determination of occlusal vertical dimension reliable? *Compend Contin Educ Dent* 18: 1214-1215, 1219-1223
- Rivera-Morales W C, Mohl N D (1991): Relationship of occlusal vertical dimension to the health of the masticatory system. *J Prosthet Dent* 65: 547-553
- Rivera-Morales W C, Mohl N D (1992): Restoration of the vertical dimension of occlusion in the severely worn dentition. *Dent Clin North Am* 36: 651-664
- Tallgren A (1972): The continuing reduction of the residual alveolar ridges in complete denture wearers: a mixed-longitudinal study covering 25 years. *J Prosthet Dent* 27: 120-132
- Timmerman R, Stoker G T, Wismeijer D, Oosterveld P, Vermeeren J I, van Waas M A (2004): An eight-year follow-up to a randomized clinical trial of participant satisfaction with three types of mandibular implant-retained overdentures. *J Dent Res* 83: 630-633
- Vig R G, Brundo G C (1978): The kinetics of anterior tooth display. *J Prosthet Dent* 39: 502-504