

| Patient | Pain medication | Week | | | | | | | | | Responder* |
|---------|--------------------|--------|--------|--------|-------|-------|-------|-------|-------|--------|------------|
| | | 1-6 | 7-12 | 13-18 | 19-24 | 25-30 | 31-36 | 37-42 | 43-48 | 49-end | |
| 73 | Diclofenac eq. | 150 | 75 | 0 | 0 | 0 | 75 | 0 | 0 | 0 | X |
| 79 | Paracetamol | 0 | 0 | 0 | 0 | 0 | 1.000 | 0 | 0 | 0 | X |
| 81 | Diclofenac eq. | 239 | | | | | | | | | |
| | Paracetamol | 500 | | | | | | | | | |
| 85 | Diclofenac eq. | 313 | | | | | | | | | |
| 86 | Diclofenac eq. | 0 | 0 | 50 | 0 | 0 | 0 | 0 | 0 | 0 | X |
| | Paracetamol | 0 | 0 | 0 | 1.000 | 0 | 0 | 0 | 0 | 0 | |
| 87 | Diclofenac eq. | 500 | 0 | 0 | | | | | | | |
| | Paracetamol | 3.000 | 0 | 0 | | | | | | | |
| 95 | Diclofenac eq. | 1.056 | 0 | 0 | 0 | 6.150 | 0 | 0 | 0 | 0 | |
| 97 | Diclofenac eq. | 0 | 200 | 0 | 0 | | | | | | |
| 98 | Diclofenac eq. | 450 | 300 | | | | | | | | |
| 101 | Paracetamol | 2.000 | 500 | 1.500 | 3.500 | 2.500 | 0 | | | | |
| 102 | Paracetamol | 29.750 | 26.000 | | | | | | | | |
| 105 | Diclofenac eq. | 3.150 | 3.150 | | | | | | | | |
| 106 | Diclofenac eq. | 0 | 900 | 0 | 0 | | | | | | |
| 114 | Diclofenac eq. | 900 | 0 | 0 | 0 | 800 | 0 | 0 | 0 | 0 | X |
| 119 | Diclofenac eq. | 375 | 225 | 75 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 130 | Diclofenac eq. | 6.150 | 6.450 | 12.900 | | | | | | | |
| | Paracetamol | 0 | 0 | 500 | | | | | | | |
| 142 | Diclofenac eq. | 2.725 | 575 | 75 | 0 | 50 | 0 | 0 | 50 | 1.050 | |
| 146 | Diclofenac eq. | 0 | 1.400 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | X |
| 147 | Central analgesics | 4.200 | 4.200 | 4.200 | 4.200 | 4.200 | 4.200 | 1.400 | 0 | 0 | X |
| 150 | Diclofenac eq. | 0 | 350 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | X |
| 156 | Diclofenac eq. | 2.400 | 350 | 650 | 1.150 | 400 | 5.000 | 4.200 | 6.500 | 3.450 | X |
| 157 | Diclofenac eq. | 1.625 | 350 | 500 | 200 | 0 | 1.050 | 300 | 0 | 0 | X |

Drop-out

Responders using analgesics during the last preceding period

* response at individual study end