

Demographic and Anamnestic data (n = 142 patients)

	Low Back		Hip		Knee		Total	
	N	(%)	N	(%)	N	(%)	N	(%)
Gender								
Male	30	(24.6)	1	(9.1)	2	(22.2)	33	(23.2)
Female	92	(75.4)	10	(90.9)	7	(77.8)	109	(76.8)
Total	122	(100)	11	(100)	9	(100)	142	(100)

	Group	N	Min	Max	P25	Median	P75	Mean	Std
Age [years]	Low Back	122	30.7	81.9	50.8	59.1	67.0	58.7	10.9
	Hip	11	52.8	80.0	60.1	68.1	69.9	66.4	7.3
	Knee	9	50.6	70.3	58.1	62.4	65.2	61.8	6.2
	Total	142	30.7	81.9	51.9	60.7	67.7	59.5	10.7
Height [cm]	Low Back	122	149.0	190.0	162.0	165.0	172.0	167.2	8.1
	Hip	11	151.0	176.0	156.0	163.0	173.0	163.9	8.6
	Knee	9	160.0	184.0	165.0	167.0	173.0	168.9	7.7
	Total	142	149.0	190.0	162.0	165.5	172.0	167.0	8.1
Body Weight [kg]	Low Back	122	48.0	128.0	65.0	76.3	84.0	76.5	15.6
	Hip	11	46.0	85.2	60.0	75.0	82.0	69.7	12.9
	Knee	9	56.0	99.0	60.0	80.0	85.0	75.8	15.4
	Total	142	46.0	128.0	64.0	76.0	84.0	75.9	15.4
Body Mass Index [kg/cm2]	Low Back	122	16.9	44.6	23.6	26.9	29.8	27.4	5.4
	Hip	11	20.2	29.7	23.0	24.8	28.6	25.8	3.1
	Knee	9	19.6	31.2	22.7	28.1	29.2	26.5	4.5
	Total	142	16.9	44.6	23.5	26.9	29.7	27.2	5.2
Duration of specific pain [months]	Low Back	122	6.0	600.0	120.0	180.0	240.0	196.4	126.1
	Hip	11	36.0	480.0	72.0	180.0	360.0	210.5	152.2
	Knee	9	18.0	360.0	24.0	120.0	300.0	154.0	136.0
	Total	142	6.0	600.0	120.0	180.0	240.0	194.8	128.3
Acute Exacerbation [weeks]	Low Back	122	2.0	960.0	12.0	20.0	48.0	63.0	118.6
	Hip	11	6.0	288.0	24.0	48.0	96.0	84.9	93.9
	Knee	9	4.0	480.0	8.0	24.0	240.0	149.3	202.4
	Total	142	2.0	960.0	12.0	24.0	56.0	70.2	124.4

Analysis Variable	P-value Kruskal-Wallis Test
Age	0.044
Height	0.388
Body Weight	0.528
Body Mass Index	0.745
Duration of specific pain	0.537
Acute Exacerbation	0.203

Analysis Variable	Fisher's Exact Test P-value (2-Tail)
Gender	0.573