

*Arhus Index**ITT-population (LOCF)**Baseline*

		<b>N</b>	<b>Min</b>	<b>Max</b>	<b>P25</b>	<b>Median</b>	<b>P75</b>	<b>Mean</b>	<b>Std</b>
<b>Current pain</b>	Low Back	39	0.0	10.0	2.0	3.1	5.2	3.8	2.5
	Knee	13	0.0	7.4	0.5	3.8	5.0	3.4	2.5
	<b>Total</b>	<b>52</b>	<b>0.0</b>	<b>10.0</b>	<b>1.8</b>	<b>3.4</b>	<b>5.1</b>	<b>3.7</b>	<b>2.5</b>
<b>Maximum pain</b>	Low Back	39	4.0	10.0	5.0	6.3	8.1	6.6	1.7
	Knee	13	5.0	10.0	5.0	6.7	7.4	6.4	1.5
	<b>Total</b>	<b>52</b>	<b>4.0</b>	<b>10.0</b>	<b>5.0</b>	<b>6.3</b>	<b>8.0</b>	<b>6.6</b>	<b>1.6</b>
<b>Average pain</b>	Low Back	39	1.0	8.0	3.0	4.2	5.4	4.3	1.7
	Knee	13	1.6	7.7	3.8	5.0	5.0	4.5	1.5
	<b>Total</b>	<b>52</b>	<b>1.0</b>	<b>8.0</b>	<b>3.1</b>	<b>4.6</b>	<b>5.3</b>	<b>4.3</b>	<b>1.6</b>
<b>Three item pain Index</b>	Low Back	39	7.0	26.8	10.5	14.0	18.7	14.7	5.0
	Knee	13	7.7	20.0	12.5	14.3	16.9	14.3	3.7
	<b>Total</b>	<b>52</b>	<b>7.0</b>	<b>26.8</b>	<b>10.9</b>	<b>14.3</b>	<b>18.2</b>	<b>14.6</b>	<b>4.7</b>
<b>Component Pain</b>	Low Back	39	7.0	48.0	10.7	15.5	27.2	20.3	11.8
	Knee								
	<b>Total</b>	<b>39</b>	<b>7.0</b>	<b>48.0</b>	<b>10.7</b>	<b>15.5</b>	<b>27.2</b>	<b>20.3</b>	<b>11.8</b>
<b>Component Invalidity</b>	Low Back	39	2.0	30.0	7.0	13.0	18.0	13.3	6.6
	Knee	13	6.0	25.0	9.0	15.0	16.0	13.6	6.0
	<b>Total</b>	<b>52</b>	<b>2.0</b>	<b>30.0</b>	<b>8.0</b>	<b>13.5</b>	<b>17.0</b>	<b>13.4</b>	<b>6.4</b>

***Arhus Index******ITT-population (LOCF)******Baseline***

<b>Analysis Variable</b>	<b>P-value Wilcoxon Test (Two-sided)</b>
Current pain	0.767
Maximum pain	0.714
Average pain	0.742
Three item pain Index	0.983
Component Invalidity	0.815

*Arhus Index**ITT-population (LOCF)**Visit 1 (6 weeks)*

		<b>N</b>	<b>Min</b>	<b>Max</b>	<b>P25</b>	<b>Median</b>	<b>P75</b>	<b>Mean</b>	<b>Std</b>
<b>Current pain</b>	Low Back	39	0.0	8.0	1.0	3.0	4.5	3.1	2.3
	Knee	13	0.0	9.0	2.0	3.0	5.0	3.5	2.6
	<b>Total</b>	<b>52</b>	<b>0.0</b>	<b>9.0</b>	<b>1.0</b>	<b>3.0</b>	<b>5.0</b>	<b>3.2</b>	<b>2.4</b>
<b>Maximum pain</b>	Low Back	39	2.0	10.0	4.0	5.9	7.0	5.8	2.1
	Knee	13	2.0	10.0	5.0	5.0	7.0	5.8	2.4
	<b>Total</b>	<b>52</b>	<b>2.0</b>	<b>10.0</b>	<b>4.5</b>	<b>5.9</b>	<b>7.0</b>	<b>5.8</b>	<b>2.2</b>
<b>Average pain</b>	Low Back	39	0.0	8.0	2.0	3.5	5.0	3.6	1.7
	Knee	13	0.0	7.7	2.5	3.6	5.1	3.8	2.1
	<b>Total</b>	<b>52</b>	<b>0.0</b>	<b>8.0</b>	<b>2.1</b>	<b>3.6</b>	<b>5.0</b>	<b>3.7</b>	<b>1.8</b>
<b>Three item pain Index</b>	Low Back	39	2.0	26.0	8.0	12.0	15.5	12.5	5.5
	Knee	13	2.0	22.0	10.0	13.6	17.1	13.1	5.7
	<b>Total</b>	<b>52</b>	<b>2.0</b>	<b>26.0</b>	<b>8.5</b>	<b>12.8</b>	<b>16.1</b>	<b>12.6</b>	<b>5.5</b>
<b>Component Pain</b>	Low Back	39	2.0	52.0	10.0	15.0	20.8	17.2	11.0
	Knee								
	<b>Total</b>	<b>39</b>	<b>2.0</b>	<b>52.0</b>	<b>10.0</b>	<b>15.0</b>	<b>20.8</b>	<b>17.2</b>	<b>11.0</b>
<b>Component Invalidity</b>	Low Back	39	1.0	24.0	6.0	11.0	15.0	10.8	6.1
	Knee	13	2.0	28.0	6.0	8.0	11.0	9.9	7.2
	<b>Total</b>	<b>52</b>	<b>1.0</b>	<b>28.0</b>	<b>6.0</b>	<b>10.0</b>	<b>14.5</b>	<b>10.6</b>	<b>6.3</b>

*Arhus Index**ITT-population (LOCF)**Visit 1 (6 weeks)*

<b>Analysis Variable</b>	<b>P-value Wilcoxon Test (Two-sided)</b>
Current pain	0.618
Maximum pain	0.899
Average pain	0.695
Three item pain Index	0.612
Component Invalidity	0.446

*Arhus Index improvement (%) from baseline**ITT-population (LOCF)**Visit 1 (6 weeks)*

		<b>N</b>	<b>Min</b>	<b>Max</b>	<b>P25</b>	<b>Median</b>	<b>P75</b>	<b>Mean</b>	<b>Std</b>
<b>Current pain [%]</b>	Low Back	36	-337.5	100.0	-17.0	14.3	64.8	6.4	89.8
	Knee	10	-300.0	100.0	-27.9	10.5	70.3	-21.3	124.7
	<b>Total</b>	<b>46</b>	<b>-337.5</b>	<b>100.0</b>	<b>-24.0</b>	<b>14.3</b>	<b>66.7</b>	<b>0.3</b>	<b>97.6</b>
<b>Three item pain Index [%]</b>	Low Back	39	-57.1	81.8	-3.7	8.1	40.8	12.6	33.3
	Knee	13	-56.0	90.0	-23.4	0.0	8.1	4.2	39.0
	<b>Total</b>	<b>52</b>	<b>-57.1</b>	<b>90.0</b>	<b>-12.6</b>	<b>6.1</b>	<b>38.0</b>	<b>10.5</b>	<b>34.6</b>
<b>Component Pain [%]</b>	Low Back	39	-57.1	88.9	-18.5	10.8	29.7	9.5	36.4
	Knee								
	<b>Total</b>	<b>39</b>	<b>-57.1</b>	<b>88.9</b>	<b>-18.5</b>	<b>10.8</b>	<b>29.7</b>	<b>9.5</b>	<b>36.4</b>
<b>Component Invalidity [%]</b>	Low Back	39	-100.0	93.8	0.0	14.3	31.3	15.5	34.0
	Knee	13	-42.9	87.5	-12.0	11.8	62.5	20.9	43.4
	<b>Total</b>	<b>52</b>	<b>-100.0</b>	<b>93.8</b>	<b>0.0</b>	<b>13.8</b>	<b>33.9</b>	<b>16.8</b>	<b>36.2</b>

***Arhus Index improvement (%) from baseline******ITT-population (LOCF)******Visit 1 (6 weeks)***

<b>Analysis Variable</b>	<b>P-value Wilcoxon Test (Two-sided)</b>
Current pain [%]	0.650
Three item pain Index [%]	0.341
Component Invalidity [%]	0.992

*Arhus Index**ITT-population (LOCF)**Visit 2 (12 weeks)*

		<b>N</b>	<b>Min</b>	<b>Max</b>	<b>P25</b>	<b>Median</b>	<b>P75</b>	<b>Mean</b>	<b>Std</b>
<b>Current pain</b>	Low Back	39	0.0	8.1	0.0	1.5	3.8	2.0	2.2
	Knee	13	0.0	6.2	0.5	1.3	4.0	2.0	2.1
	<b>Total</b>	<b>52</b>	<b>0.0</b>	<b>8.1</b>	<b>0.0</b>	<b>1.4</b>	<b>3.9</b>	<b>2.0</b>	<b>2.2</b>
<b>Maximum pain</b>	Low Back	39	0.0	9.1	2.0	5.0	6.3	4.3	2.8
	Knee	13	2.0	10.0	3.0	4.4	8.0	5.5	2.9
	<b>Total</b>	<b>52</b>	<b>0.0</b>	<b>10.0</b>	<b>2.6</b>	<b>4.8</b>	<b>6.8</b>	<b>4.6</b>	<b>2.8</b>
<b>Average pain</b>	Low Back	39	0.0	7.1	1.4	2.5	3.8	2.7	1.8
	Knee	13	1.2	7.7	1.4	2.3	4.0	3.1	2.0
	<b>Total</b>	<b>52</b>	<b>0.0</b>	<b>7.7</b>	<b>1.4</b>	<b>2.5</b>	<b>4.0</b>	<b>2.8</b>	<b>1.9</b>
<b>Three item pain Index</b>	Low Back	39	0.0	20.5	4.1	8.7	14.0	9.0	6.2
	Knee	13	4.5	18.5	5.6	10.0	14.5	10.6	5.3
	<b>Total</b>	<b>52</b>	<b>0.0</b>	<b>20.5</b>	<b>5.3</b>	<b>8.9</b>	<b>14.4</b>	<b>9.4</b>	<b>6.0</b>
<b>Component Pain</b>	Low Back	39	0.0	43.5	5.0	13.5	21.0	13.7	11.8
	Knee								
	<b>Total</b>	<b>39</b>	<b>0.0</b>	<b>43.5</b>	<b>5.0</b>	<b>13.5</b>	<b>21.0</b>	<b>13.7</b>	<b>11.8</b>
<b>Component Invalidity</b>	Low Back	39	0.0	30.0	4.0	9.0	12.0	9.4	6.7
	Knee	13	3.0	14.0	5.0	7.0	9.0	7.4	3.1
	<b>Total</b>	<b>52</b>	<b>0.0</b>	<b>30.0</b>	<b>4.0</b>	<b>8.0</b>	<b>12.0</b>	<b>8.9</b>	<b>6.0</b>

***Arhus Index******ITT-population (LOCF)******Visit 2 (12 weeks)***

<b>Analysis Variable</b>	<b>P-value Wilcoxon Test (Two-sided)</b>
Current pain	0.746
Maximum pain	0.300
Average pain	0.688
Three item pain Index	0.499
Component Invalidity	0.433



*Arhus Index improvement (%) from baseline**ITT-population (LOCF)**Visit 2 (12 weeks)*

		<b>N</b>	<b>Min</b>	<b>Max</b>	<b>P25</b>	<b>Median</b>	<b>P75</b>	<b>Mean</b>	<b>Std</b>
<b>Current pain [%]</b>	Low Back	36	-200.0	100.0	8.3	55.9	100.0	41.0	67.8
	Knee	10	-24.2	100.0	0.0	68.2	86.8	46.1	48.5
	<b>Total</b>	<b>46</b>	<b>-200.0</b>	<b>100.0</b>	<b>0.0</b>	<b>60.7</b>	<b>100.0</b>	<b>42.1</b>	<b>63.6</b>
<b>Three item pain Index [%]</b>	Low Back	39	-71.4	100.0	4.8	38.4	75.0	35.3	48.5
	Knee	13	-48.0	71.6	4.7	23.5	51.7	24.8	34.5
	<b>Total</b>	<b>52</b>	<b>-71.4</b>	<b>100.0</b>	<b>4.7</b>	<b>35.3</b>	<b>66.1</b>	<b>32.7</b>	<b>45.4</b>
<b>Component Pain [%]</b>	Low Back	39	-230.0	100.0	0.0	33.7	80.8	21.1	75.1
	Knee								
	<b>Total</b>	<b>39</b>	<b>-230.0</b>	<b>100.0</b>	<b>0.0</b>	<b>33.7</b>	<b>80.8</b>	<b>21.1</b>	<b>75.1</b>
<b>Component Invalidity [%]</b>	Low Back	39	-100.0	100.0	0.0	28.6	52.9	26.9	43.3
	Knee	13	0.0	75.0	17.6	44.4	60.0	39.8	25.8
	<b>Total</b>	<b>52</b>	<b>-100.0</b>	<b>100.0</b>	<b>3.6</b>	<b>31.0</b>	<b>55.9</b>	<b>30.1</b>	<b>39.8</b>

***Arhus Index improvement (%) from baseline******ITT-population (LOCF)******Visit 2 (12 weeks)***

<b>Analysis Variable</b>	<b>P-value Wilcoxon Test (Two-sided)</b>
Current pain [%]	0.686
Three item pain Index [%]	0.357
Component Invalidity [%]	0.315