

*Weekly dose**ITT-population (LOCF)*

	Group	N	Min	Max	P25	Median	P75	Mean	Std
<b>Week 1</b>	Low Back	38	70	168	105.0	126.0	168.0	128.7	34.7
	Knee	13	105	168	126.0	140.0	150.0	136.4	23.1
	<b>Total</b>	<b>51</b>	<b>70</b>	<b>168</b>	<b>105.0</b>	<b>126.0</b>	<b>168.0</b>	<b>130.7</b>	<b>32.1</b>
<b>Week 2</b>	Low Back	38	40	168	105.0	126.0	168.0	126.7	38.3
	Knee	13	105	350	126.0	140.0	168.0	151.3	63.9
	<b>Total</b>	<b>51</b>	<b>40</b>	<b>350</b>	<b>105.0</b>	<b>126.0</b>	<b>168.0</b>	<b>133.0</b>	<b>46.7</b>
<b>Week 3</b>	Low Back	38	30	168	105.0	126.0	168.0	126.6	42.1
	Knee	13	70	350	110.0	126.0	168.0	146.3	67.6
	<b>Total</b>	<b>51</b>	<b>30</b>	<b>350</b>	<b>105.0</b>	<b>126.0</b>	<b>168.0</b>	<b>131.6</b>	<b>49.8</b>
<b>Week 4</b>	Low Back	38	35	168	105.0	126.0	168.0	128.6	39.1
	Knee	13	70	350	105.0	126.0	168.0	143.2	70.2
	<b>Total</b>	<b>51</b>	<b>35</b>	<b>350</b>	<b>105.0</b>	<b>126.0</b>	<b>168.0</b>	<b>132.3</b>	<b>48.5</b>
<b>Week 5</b>	Low Back	38	35	192	105.0	126.0	168.0	128.5	42.3
	Knee	13	70	350	126.0	126.0	168.0	147.5	67.0
	<b>Total</b>	<b>51</b>	<b>35</b>	<b>350</b>	<b>105.0</b>	<b>126.0</b>	<b>168.0</b>	<b>133.3</b>	<b>49.7</b>
<b>Week 6</b>	Low Back	38	35	252	105.0	126.0	168.0	129.5	46.2
	Knee	13	84	350	126.0	126.0	168.0	148.2	66.1
	<b>Total</b>	<b>51</b>	<b>35</b>	<b>350</b>	<b>105.0</b>	<b>126.0</b>	<b>168.0</b>	<b>134.3</b>	<b>51.9</b>
<b>Week 7</b>	Low Back	38	35	252	105.0	140.0	168.0	134.5	45.0
	Knee	13	126	350	126.0	140.0	168.0	160.5	60.4
	<b>Total</b>	<b>51</b>	<b>35</b>	<b>350</b>	<b>126.0</b>	<b>140.0</b>	<b>168.0</b>	<b>141.1</b>	<b>50.0</b>
<b>Week 8</b>	Low Back	38	35	252	126.0	140.0	168.0	136.1	44.7
	Knee	13	126	350	126.0	168.0	168.0	163.7	59.5
	<b>Total</b>	<b>51</b>	<b>35</b>	<b>350</b>	<b>126.0</b>	<b>140.0</b>	<b>168.0</b>	<b>143.2</b>	<b>49.8</b>
<b>Week 9</b>	Low Back	38	35	252	126.0	140.0	168.0	136.4	44.7
	Knee	13	126	350	126.0	140.0	168.0	160.5	60.4
	<b>Total</b>	<b>51</b>	<b>35</b>	<b>350</b>	<b>126.0</b>	<b>140.0</b>	<b>168.0</b>	<b>142.6</b>	<b>49.7</b>
<b>Week 10</b>	Low Back	38	35	252	126.0	133.0	168.0	134.9	44.3
	Knee	13	126	350	126.0	168.0	168.0	166.9	58.4
	<b>Total</b>	<b>51</b>	<b>35</b>	<b>350</b>	<b>126.0</b>	<b>140.0</b>	<b>168.0</b>	<b>143.0</b>	<b>49.7</b>
<b>Week 11</b>	Low Back	38	35	252	126.0	133.0	168.0	135.0	44.4
	Knee	13	126	350	126.0	168.0	168.0	166.9	58.4
	<b>Total</b>	<b>51</b>	<b>35</b>	<b>350</b>	<b>126.0</b>	<b>140.0</b>	<b>168.0</b>	<b>143.2</b>	<b>49.8</b>

<b>Group</b>		<b>N</b>	<b>Min</b>	<b>Max</b>	<b>P25</b>	<b>Median</b>	<b>P75</b>	<b>Mean</b>	<b>Std</b>
<b>Week 12</b>	Low Back	38	35	252	126.0	133.0	168.0	135.0	44.4
	Knee	13	126	350	126.0	168.0	168.0	166.9	58.4
	<b>Total</b>	<b>51</b>	<b>35</b>	<b>350</b>	<b>126.0</b>	<b>140.0</b>	<b>168.0</b>	<b>143.2</b>	<b>49.8</b>

*Weekly dose**ITT-population (LOCF)*

<b>Analysis Variable</b>	<b>P-value Wilcoxon Test (Two-sided)</b>
Week 1	0.573
Week 2	0.345
Week 3	0.730
Week 4	0.902
Week 5	0.747
Week 6	0.689
Week 7	0.348
Week 8	0.271
Week 9	0.494
Week 10	0.096
Week 11	0.111
Week 12	0.111

***Dose reduction (%) from Week 1***  
***Of note: negative values represent dose increase***  
***ITT-population (LOCF)***

	Group	N	Min	Max	P25	Median	P75	Mean	Std
<b>Week 2</b>	Low Back	38	-71	50	0.0	0.0	0.0	0.7	23.1
	Knee	13	-133	5	0.0	0.0	0.0	-9.9	37.1
	<b>Total</b>	<b>51</b>	<b>-133</b>	<b>50</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>-2.0</b>	<b>27.3</b>
<b>Week 3</b>	Low Back	38	-100	60	0.0	0.0	0.0	2.0	26.3
	Knee	13	-133	33	0.0	0.0	0.0	-5.7	39.7
	<b>Total</b>	<b>51</b>	<b>-133</b>	<b>60</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>30.1</b>
<b>Week 4</b>	Low Back	38	-100	55	0.0	0.0	0.0	-0.7	25.4
	Knee	13	-133	50	0.0	0.0	0.0	-3.5	42.1
	<b>Total</b>	<b>51</b>	<b>-133</b>	<b>55</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>-1.4</b>	<b>30.0</b>
<b>Week 5</b>	Low Back	38	-100	55	0.0	0.0	0.0	-0.2	27.2
	Knee	13	-133	33	0.0	0.0	0.0	-6.6	39.2
	<b>Total</b>	<b>51</b>	<b>-133</b>	<b>55</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>-1.8</b>	<b>30.4</b>
<b>Week 6</b>	Low Back	38	-100	55	0.0	0.0	0.0	-0.6	28.5
	Knee	13	-133	20	0.0	0.0	4.5	-7.2	38.5
	<b>Total</b>	<b>51</b>	<b>-133</b>	<b>55</b>	<b>0.0</b>	<b>0.0</b>	<b>0.0</b>	<b>-2.3</b>	<b>31.1</b>
<b>Week 7</b>	Low Back	38	-140	55	-16.7	0.0	0.0	-6.6	37.1
	Knee	13	-133	10	-20.0	0.0	0.0	-18.4	39.1
	<b>Total</b>	<b>51</b>	<b>-140</b>	<b>55</b>	<b>-20.0</b>	<b>0.0</b>	<b>0.0</b>	<b>-9.6</b>	<b>37.6</b>
<b>Week 8</b>	Low Back	38	-140	55	-20.0	0.0	0.0	-8.3	38.1
	Knee	13	-133	10	-20.0	0.0	0.0	-21.4	40.7
	<b>Total</b>	<b>51</b>	<b>-140</b>	<b>55</b>	<b>-20.0</b>	<b>0.0</b>	<b>0.0</b>	<b>-11.6</b>	<b>38.8</b>
<b>Week 9</b>	Low Back	38	-140	55	-31.4	0.0	0.0	-8.6	38.3
	Knee	13	-133	10	-20.0	0.0	0.0	-18.4	39.1
	<b>Total</b>	<b>51</b>	<b>-140</b>	<b>55</b>	<b>-20.0</b>	<b>0.0</b>	<b>0.0</b>	<b>-11.1</b>	<b>38.3</b>
<b>Week 10</b>	Low Back	38	-140	55	-33.3	0.0	12.5	-7.6	39.1
	Knee	13	-133	10	-60.0	0.0	0.0	-24.5	42.1
	<b>Total</b>	<b>51</b>	<b>-140</b>	<b>55</b>	<b>-33.3</b>	<b>0.0</b>	<b>0.0</b>	<b>-11.9</b>	<b>40.1</b>
<b>Week 11</b>	Low Back	38	-140	55	-16.7	0.0	0.0	-7.4	37.9
	Knee	13	-133	10	-60.0	0.0	0.0	-24.5	42.1
	<b>Total</b>	<b>51</b>	<b>-140</b>	<b>55</b>	<b>-20.0</b>	<b>0.0</b>	<b>0.0</b>	<b>-11.8</b>	<b>39.3</b>
<b>Week 12</b>	Low Back	38	-140	55	-16.7	0.0	0.0	-7.4	37.9
	Knee	13	-133	10	-60.0	0.0	0.0	-24.5	42.1
	<b>Total</b>	<b>51</b>	<b>-140</b>	<b>55</b>	<b>-20.0</b>	<b>0.0</b>	<b>0.0</b>	<b>-11.8</b>	<b>39.3</b>

***Dose reduction (%) from Week 1******ITT-population (LOCF)***

<b>Analysis Variable</b>	<b>P-value Wilcoxon Test (Two-sided)</b>
Week 2	0.896
Week 3	0.668
Week 4	0.398
Week 5	0.567
Week 6	0.738
Week 7	0.392
Week 8	0.417
Week 9	0.537
Week 10	0.185
Week 11	0.284
Week 12	0.284