

*Health Assessment Questionnaire (HAQ)**ITT-population (LOCF)*

	Group	N	Min	Max	P25	Median	P75	Mean	Std
Baseline	Low Back	39	0.0	33.0	4.0	9.0	13.0	9.6	7.6
	Knee	13	0.0	24.0	4.0	8.0	16.0	9.8	7.8
	Total	52	0.0	33.0	4.0	8.5	13.5	9.7	7.6
Visit 1 (6 weeks)	Low Back	39	0.0	39.0	2.0	8.0	12.0	8.5	8.2
	Knee	13	0.0	18.0	4.0	6.0	11.0	7.5	5.9
	Total	52	0.0	39.0	2.0	7.5	11.5	8.2	7.6
Visit 2 (12 weeks)	Low Back	39	0.0	39.0	1.0	7.0	10.0	8.0	8.5
	Knee	13	0.0	17.0	2.0	6.0	9.0	6.5	5.5
	Total	52	0.0	39.0	2.0	6.5	10.0	7.6	7.8

Health Assessment Questionnaire (HAQ)***ITT-population (LOCF)***

Analysis Variable	P-value Kruskal-Wallis Test
Baseline	0.975
Visit 1 (6 weeks)	0.975
Visit 2 (12 weeks)	0.924

*HAQ Improvement (%) from baseline**ITT-population (LOCF)*

	Group	N	Min	Max	P25	Median	P75	Mean	Std
Visit 1 (6 weeks)	Low Back	37	-100.0	100.0	0.0	20.0	42.9	20.6	45.7
	Knee	12	-60.0	75.0	-10.0	23.4	33.3	12.6	37.1
	Total	49	-100.0	100.0	0.0	21.4	38.9	18.7	43.5
Visit 2 (12 weeks)	Low Back	37	-100.0	100.0	-18.2	30.8	66.7	22.6	55.2
	Knee	12	-20.0	72.7	5.0	33.0	61.5	32.1	31.9
	Total	49	-100.0	100.0	0.0	31.3	66.7	24.9	50.4

HAQ Improvement (%) from baseline***ITT-population (LOCF)***

Analysis Variable	P-value Kruskal-Wallis Test
Visit 1 (6 weeks)	0.797
Visit 2 (12 weeks)	0.780