

*Arhus Index**mPP-population**Baseline*

		<b>N</b>	<b>Min</b>	<b>Max</b>	<b>P25</b>	<b>Median</b>	<b>P75</b>	<b>Mean</b>	<b>Std</b>
<b>Current pain</b>	Low Back	39	0.0	10.0	2.0	3.1	5.2	3.8	2.5
	Knee	13	0.0	7.4	0.5	3.8	5.0	3.4	2.5
	<b>Total</b>	<b>52</b>	<b>0.0</b>	<b>10.0</b>	<b>1.8</b>	<b>3.4</b>	<b>5.1</b>	<b>3.7</b>	<b>2.5</b>
<b>Maximum pain</b>	Low Back	39	4.0	10.0	5.0	6.3	8.1	6.6	1.7
	Knee	13	5.0	10.0	5.0	6.7	7.4	6.4	1.5
	<b>Total</b>	<b>52</b>	<b>4.0</b>	<b>10.0</b>	<b>5.0</b>	<b>6.3</b>	<b>8.0</b>	<b>6.6</b>	<b>1.6</b>
<b>Average pain</b>	Low Back	39	1.0	8.0	3.0	4.2	5.4	4.3	1.7
	Knee	13	1.6	7.7	3.8	5.0	5.0	4.5	1.5
	<b>Total</b>	<b>52</b>	<b>1.0</b>	<b>8.0</b>	<b>3.1</b>	<b>4.6</b>	<b>5.3</b>	<b>4.3</b>	<b>1.6</b>
<b>Three item pain Index</b>	Low Back	39	7.0	26.8	10.5	14.0	18.7	14.7	5.0
	Knee	13	7.7	20.0	12.5	14.3	16.9	14.3	3.7
	<b>Total</b>	<b>52</b>	<b>7.0</b>	<b>26.8</b>	<b>10.9</b>	<b>14.3</b>	<b>18.2</b>	<b>14.6</b>	<b>4.7</b>
<b>Component Pain</b>	Low Back	39	7.0	48.0	10.7	15.5	27.2	20.3	11.8
	Knee								.
	<b>Total</b>	<b>39</b>	<b>7.0</b>	<b>48.0</b>	<b>10.7</b>	<b>15.5</b>	<b>27.2</b>	<b>20.3</b>	<b>11.8</b>
<b>Component Invalidity</b>	Low Back	39	2.0	30.0	7.0	13.0	18.0	13.3	6.6
	Knee	13	6.0	25.0	9.0	15.0	16.0	13.6	6.0
	<b>Total</b>	<b>52</b>	<b>2.0</b>	<b>30.0</b>	<b>8.0</b>	<b>13.5</b>	<b>17.0</b>	<b>13.4</b>	<b>6.4</b>

***Arhus Index******mPP-population******Baseline***

<b>Analysis Variable</b>	<b>P-value Wilcoxon Test (Two-sided)</b>
Current pain	0.767
Maximum pain	0.714
Average pain	0.742
Three item pain Index	0.983
Component Invalidity	0.815

*Arhus Index**mPP-population**Visit 1 (6 weeks)*

		<b>N</b>	<b>Min</b>	<b>Max</b>	<b>P25</b>	<b>Median</b>	<b>P75</b>	<b>Mean</b>	<b>Std</b>
<b>Current pain</b>	Low Back	34	0.0	8.0	1.0	2.6	4.5	3.0	2.4
	Knee	12	0.0	9.0	2.1	3.5	5.3	3.8	2.5
	<b>Total</b>	<b>46</b>	<b>0.0</b>	<b>9.0</b>	<b>1.0</b>	<b>3.0</b>	<b>5.0</b>	<b>3.2</b>	<b>2.4</b>
<b>Maximum pain</b>	Low Back	34	2.0	10.0	4.0	5.3	7.0	5.7	2.2
	Knee	12	2.0	9.8	4.5	5.0	6.8	5.4	2.2
	<b>Total</b>	<b>46</b>	<b>2.0</b>	<b>10.0</b>	<b>4.0</b>	<b>5.0</b>	<b>7.0</b>	<b>5.6</b>	<b>2.2</b>
<b>Average pain</b>	Low Back	34	0.0	8.0	2.0	3.0	4.4	3.4	1.8
	Knee	12	0.0	6.2	2.4	3.3	4.9	3.5	1.8
	<b>Total</b>	<b>46</b>	<b>0.0</b>	<b>8.0</b>	<b>2.0</b>	<b>3.0</b>	<b>4.7</b>	<b>3.4</b>	<b>1.8</b>
<b>Three item pain Index</b>	Low Back	34	2.0	26.0	7.1	11.1	15.5	12.1	5.7
	Knee	12	2.0	22.0	9.8	13.6	16.9	12.7	5.7
	<b>Total</b>	<b>46</b>	<b>2.0</b>	<b>26.0</b>	<b>8.0</b>	<b>11.6</b>	<b>15.5</b>	<b>12.2</b>	<b>5.6</b>
<b>Component Pain</b>	Low Back	34	2.0	52.0	9.8	14.5	20.8	17.1	11.3
	Knee								
	<b>Total</b>	<b>34</b>	<b>2.0</b>	<b>52.0</b>	<b>9.8</b>	<b>14.5</b>	<b>20.8</b>	<b>17.1</b>	<b>11.3</b>
<b>Component Invalidity</b>	Low Back	34	1.0	24.0	6.0	11.0	16.0	11.1	6.3
	Knee	12	2.0	28.0	5.0	7.5	13.0	10.0	7.5
	<b>Total</b>	<b>46</b>	<b>1.0</b>	<b>28.0</b>	<b>6.0</b>	<b>10.0</b>	<b>15.0</b>	<b>10.8</b>	<b>6.5</b>

***Arhus Index******mPP-population******Visit 1 (6 weeks)***

<b>Analysis Variable</b>	<b>P-value Wilcoxon Test (Two-sided)</b>
Current pain	0.303
Maximum pain	0.762
Average pain	0.754
Three item pain Index	0.617
Component Invalidity	0.460

*Arhus Index improvement (%) from baseline**mPP-population**Visit 1 (6 weeks)*

		<b>N</b>	<b>Min</b>	<b>Max</b>	<b>P25</b>	<b>Median</b>	<b>P75</b>	<b>Mean</b>	<b>Std</b>
<b>Current pain [%]</b>	Low Back	31	-337.5	100.0	-24.0	20.0	86.0	7.4	97.0
	Knee	10	-300.0	100.0	-27.9	10.5	70.3	-21.3	124.7
	<b>Total</b>	<b>41</b>	<b>-337.5</b>	<b>100.0</b>	<b>-24.0</b>	<b>20.0</b>	<b>70.3</b>	<b>0.4</b>	<b>103.5</b>
<b>Three item pain Index [%]</b>	Low Back	34	-57.1	81.8	-12.5	20.0	41.4	14.5	35.3
	Knee	12	-56.0	90.0	-23.4	-1.6	21.5	4.6	40.7
	<b>Total</b>	<b>46</b>	<b>-57.1</b>	<b>90.0</b>	<b>-16.8</b>	<b>8.1</b>	<b>40.8</b>	<b>11.9</b>	<b>36.6</b>
<b>Component Pain [%]</b>	Low Back	34	-57.1	88.9	-18.5	17.4	35.0	10.9	38.9
	Knee								.
	<b>Total</b>	<b>34</b>	<b>-57.1</b>	<b>88.9</b>	<b>-18.5</b>	<b>17.4</b>	<b>35.0</b>	<b>10.9</b>	<b>38.9</b>
<b>Component Invalidity [%]</b>	Low Back	34	-100.0	93.8	0.0	16.2	34.5	17.7	36.0
	Knee	12	-42.9	87.5	-17.1	20.9	66.0	22.7	44.9
	<b>Total</b>	<b>46</b>	<b>-100.0</b>	<b>93.8</b>	<b>0.0</b>	<b>16.2</b>	<b>37.5</b>	<b>19.0</b>	<b>38.0</b>

***Arhus Index improvement (%) from baseline******mPP-population******Visit 1 (6 weeks)***

<b>Analysis Variable</b>	<b>P-value Wilcoxon Test (Two-sided)</b>
Current pain [%]	0.552
Three item pain Index [%]	0.323
Component Invalidity [%]	0.960

*Arhus Index**mPP-population**Visit 2 (12 weeks)*

		<b>N</b>	<b>Min</b>	<b>Max</b>	<b>P25</b>	<b>Median</b>	<b>P75</b>	<b>Mean</b>	<b>Std</b>
<b>Current pain</b>	Low Back	36	0.0	8.1	0.0	1.0	3.0	1.8	2.2
	Knee	12	0.0	6.2	0.5	1.4	4.1	2.1	2.2
	<b>Total</b>	<b>48</b>	<b>0.0</b>	<b>8.1</b>	<b>0.0</b>	<b>1.2</b>	<b>3.4</b>	<b>1.9</b>	<b>2.2</b>
<b>Maximum pain</b>	Low Back	36	0.0	9.1	2.0	4.6	6.1	4.2	2.8
	Knee	12	2.0	10.0	2.8	4.2	7.6	5.2	2.7
	<b>Total</b>	<b>48</b>	<b>0.0</b>	<b>10.0</b>	<b>2.5</b>	<b>4.5</b>	<b>6.4</b>	<b>4.4</b>	<b>2.8</b>
<b>Average pain</b>	Low Back	36	0.0	7.1	1.2	2.5	3.3	2.5	1.8
	Knee	12	1.2	5.4	1.4	2.1	4.0	2.7	1.5
	<b>Total</b>	<b>48</b>	<b>0.0</b>	<b>7.1</b>	<b>1.4</b>	<b>2.5</b>	<b>3.9</b>	<b>2.6</b>	<b>1.7</b>
<b>Three item pain Index</b>	Low Back	36	0.0	20.5	3.6	7.5	13.4	8.5	6.2
	Knee	12	4.5	18.5	5.6	8.7	14.4	10.0	5.0
	<b>Total</b>	<b>48</b>	<b>0.0</b>	<b>20.5</b>	<b>4.9</b>	<b>7.5</b>	<b>13.9</b>	<b>8.9</b>	<b>5.9</b>
<b>Component Pain</b>	Low Back	36	0.0	43.5	4.6	10.4	21.1	13.6	12.3
	Knee								
	<b>Total</b>	<b>36</b>	<b>0.0</b>	<b>43.5</b>	<b>4.6</b>	<b>10.4</b>	<b>21.1</b>	<b>13.6</b>	<b>12.3</b>
<b>Component Invalidity</b>	Low Back	36	0.0	30.0	4.0	9.5	12.5	9.5	6.9
	Knee	12	3.0	14.0	4.5	7.0	9.0	7.3	3.2
	<b>Total</b>	<b>48</b>	<b>0.0</b>	<b>30.0</b>	<b>4.0</b>	<b>8.0</b>	<b>12.0</b>	<b>9.0</b>	<b>6.2</b>

***Arhus Index******mPP-population******Visit 2 (12 weeks)***

<b>Analysis Variable</b>	<b>P-value Wilcoxon Test (Two-sided)</b>
Current pain	0.394
Maximum pain	0.411
Average pain	0.793
Three item pain Index	0.543
Component Invalidity	0.403



*Arhus Index improvement (%) from baseline**mPP-population**Visit 2 (12 weeks)*

		<b>N</b>	<b>Min</b>	<b>Max</b>	<b>P25</b>	<b>Median</b>	<b>P75</b>	<b>Mean</b>	<b>Std</b>
<b>Current pain [%]</b>	Low Back	33	-200.0	100.0	25.0	63.5	100.0	44.8	69.7
	Knee	10	-24.2	100.0	0.0	68.2	86.8	46.1	48.5
	<b>Total</b>	<b>43</b>	<b>-200.0</b>	<b>100.0</b>	<b>20.0</b>	<b>66.7</b>	<b>100.0</b>	<b>45.1</b>	<b>64.8</b>
<b>Three item pain Index [%]</b>	Low Back	36	-71.4	100.0	14.3	40.8	76.8	38.2	49.4
	Knee	12	-48.0	71.6	12.3	26.8	56.3	26.9	35.2
	<b>Total</b>	<b>48</b>	<b>-71.4</b>	<b>100.0</b>	<b>14.3</b>	<b>39.4</b>	<b>70.2</b>	<b>35.4</b>	<b>46.2</b>
<b>Component Pain [%]</b>	Low Back	36	-230.0	100.0	-8.3	43.3	82.6	22.8	78.0
	Knee								
	<b>Total</b>	<b>36</b>	<b>-230.0</b>	<b>100.0</b>	<b>-8.3</b>	<b>43.3</b>	<b>82.6</b>	<b>22.8</b>	<b>78.0</b>
<b>Component Invalidity [%]</b>	Low Back	36	-100.0	100.0	3.6	34.8	54.2	29.1	44.4
	Knee	12	12.5	75.0	18.8	47.2	62.6	43.1	23.8
	<b>Total</b>	<b>48</b>	<b>-100.0</b>	<b>100.0</b>	<b>13.4</b>	<b>37.1</b>	<b>58.1</b>	<b>32.6</b>	<b>40.4</b>

***Arhus Index improvement (%) from baseline******mPP-population******Visit 2 (12 weeks)***

<b>Analysis Variable</b>	<b>P-value Wilcoxon Test (Two-sided)</b>
Current pain [%]	0.512
Three item pain Index [%]	0.311
Component Invalidity [%]	0.341